

Image Centering with a Maximum Likelihood Estimator: Application to Infrared Astronomical imaging

Damien Gratadour^{1,2}, Laurent M. Mugnier¹, Daniel Rouan²

¹ *DOTA / ONERA, BP 72, 92322 Châtillon cedex, France*

² *LESIA / Observatoire de Paris-Meudon, 92195 Meudon, France*

Abstract. We present a new, Maximum Likelihood (ML) based, method for the estimation of the shift between two images. It notably outperforms the classical cross-correlation method especially in the case of low photon levels. Moreover, it is arbitrarily subpixel, without any resampling of the image, through the maximisation of a criterion. The method was tested with simulations and was applied to the case of infrared astronomical imaging where the signal is usually very weak. We have also extended our method to the joint estimation of the shifts in a sequence of N images, and preliminary results are presented in last section.

1. Introduction

An accurate centering of a sequence of images is mandatory in thermal IR astrophysics to increase SNR while preserving the resolution of an instrument. A classical method to estimate the translation parameters is the linear cross-correlation of noisy images with a reference (Vanderlugt 1964, Kumar et al. 1992). Downie and Walkup (1994) showed that taking into account the noise statistic can greatly improve the accuracy. Carfantan & Rougé have studied the case of the subpixel estimation of the maximum of the intercorrelation of two images with various interpolation for a stationnary gaussian noise (Carfantan and Rougé 2001). Finally Guillaume et al. have studied the pixel accurate shift estimation problem in the case of poissonian noise at low photon level (Guillaume et al. 1998). We have developped a new, Maximum Likelihood (ML) based, method for the estimation of the shift between two images. It is arbitrarily subpixel, without any resampling of the image, through the maximisation of a criterion. We describe in section 2 the theoretical basis of the method. Then in section 3, we present the results obtained with simulated images for different types of noise (pure gaussian additive or mixture of stationnary gaussian and poissonian noise). Results on real data are presented in section 4. We have also extended our method to the joint estimation of the shifts in a sequence of N images, and preliminary results are presented in last section.

2. Description of the method

If we assume a reference r , the intensity at pixel (k, l) of the observed translated image i_1 can be written as:

$$i_1(k, l) = [r(x, y) * \delta(x - x_1, y - y_1)]_{\text{III}}(k, l) + b_1(k, l) \quad (1)$$

where (x_1, y_1) are the translation parameters, b is an additive noise, and III is the sampling operator. If the image is Nyquist sampled, one can reconstruct, via the Fourier domain, a shifted version of the image for any subpixel shift. If we approximate the noise in the image, i.e. a mixture of gaussian (detector) and poissonian noise, as a non-stationary gaussian noise, then the anti log-likelihood of observing an intensity $i_1(k, l)$ for the reference intensity $r(x, y)$ and for the hypothesis $\mu = (x_1, y_1)$ is given by:

$$\mathcal{J}(x_1, y_1) = \sum_{k, l} \frac{1}{2\sigma_1^2(k, l)} |i_1(k, l) - [r(x, y) * \delta(x - x_1, y - y_1)]_{\text{III}}(k, l)|^2 \quad (2)$$

where σ_1^2 is the noise variance which can be directly estimated on the image. It is easy to show that, following the two hypothesis of stationnarity of the noise and of periodicity of the reference, the ML estimate of the translation between the two images is the maximum of the linear cross-correlation of the images. When the reference is not known, one has to consider a noisy frame as a reference.

$$i_1(k, l) = [i_0(x, y) * \delta(x - x_1, y - y_1)]_{\text{III}} + b(k, l) \quad (3)$$

Where b includes both the noise in the image used as a reference and the noise in the image to be recentered. Then the anti log-likelihood to be minimize has the same expression as in equation 2 changing $r(x, y)$ into $i_0(x, y)$ and $\sigma_1^2(k, l)$ into $\sigma^2(k, l) = \sigma_1^2(k, l) + \sigma_0^2(x, y) * \delta(x - x_1, y - y_1) = 2\sigma_1^2(k, l)$:

$$\mathcal{J}(x_1, y_1) = \sum_{k, l} \frac{1}{4\sigma_1^2(k, l)} |i_1(x, y) - [i_0(x - x_1, y - y_1)]_{\text{III}}|^2 \quad (4)$$

To find the minimum of this criterion, we used a gradient type adaptive step minimization algorithm, issued from a collaboration of our team with the Groupe des Problemes Inverses at Laboratoire des Signaux et Systemes (GPI 1997). However, one has to notice that the criterion, in the case of unknown reference and considering the real noise variance contains a lot of local minima. This make the minimization difficult and so should decrease the performance of the method in this case.

3. Results with simulation

The method has been tested with simulations in the case of a mixture of gaussian (detector) and poissonian noise. The gaussian noise variance is constant (10) and the photon level in the images ranges from 1 to 10^6 . The cross-correlation of the two images is interpolated around its maximum to provide a sub-pixel estimation. In the case of the known reference, our method, considering a constant




Figure 1. Adaptive optics image of ARP 220 in the L-band with NACO at VLT. Left image, frames registered with a classical cross-correlation method and averaged, right, frames registered with our algorithm and averaged..

noise variance, outperforms the cross-correlation at very low photon level (i.e. number of photons smaller or equal to variance of the detector noise). When we consider the real noise variance, our method gives more accurate results and slightly outperforms the cross-correlation at high photon levels.

In the case of an unknown reference, the performance of our method is quite identical considering or not the real noise distribution since the criterion contains a lot of local minima in the first case. It notably outperforms the cross-correlation at low photon level and allows subpixel accuracy as soon as the number of photon per pixel is greater than the variance of the detector noise as the accuracy of the interpolated cross correlation is worst than the pixel.

4. Results with real data

The method has also been tested and used with a set of raw images of Arp 220 from NAOS-CONICA (NACO) at VLT. Arp 220 is a typical Ultra Luminous Infrared Galaxy, characterised by a very powerfull emission in infrared bands but very faint counterpart at the visible wavelengths. NACO is the only adaptive optics system that allows to servo infrared source and so achieve diffraction limited images at a large telescope of such galaxies. A series of 85 images of this galaxy have been aquired in the L-band in March 2003. The background dynamics of each image is around 80000 photons per pixel and the source dynamic at the maximum is around 200 photons per pixel. This is the case where the classical correlation of images is inefficient. Our method allows to recenter each frame with a subpixel accuracy, and so to obtain the image displayed on Fig. 1. The resolution of this image on the sky is about $0.1''$, i.e. diffraction limited for a 8-m telescope in the L-band.

This allows to compare this image to the one obtain with the space telescope in other bands giving insightfull astophysical interpretations (see Gratadour et al. 2003).

5. Joint estimation of the reference and the translation parameters

If we consider now a series of images $\{i_j(k, l)\}$ randomly shifted, and if we try to find simultaneously the shift parameters $\{\mu_j\} = \{(x_j, y_j)\}$ and the reference image $r(x, y)$, then the anti log-likelihood can be written as:

$$\mathcal{J}(\{i_j(k, l)\}; r(x, y), \{\mu_j\}) = \sum_m \sum_{k, l} \frac{1}{2\sigma_1^2(k, l)} |i_m(k, l) - [r * \delta(x - x_m, y - y_m)]_{\text{III}}(k, l)|^2$$

One can show that minimizing $\mathcal{J}(\{i_j(k, l)\}; r(x, y), \{\mu_j\})$ on $r(x, y)$ and $\{\mu_j\}$ is equivalent to minimize: $\mathcal{J}(\{i_j(k, l)\}; r(x, y) = r_{ML}(k, l), \{\mu_j\})$ on $\{\mu_j\}$, with:

$$r_{ML}(k, l) = \sum_m i_m(k, l) * \delta(x + x_m, y + y_m) \quad (5)$$

It can additionally be shown (Blanc et al. 2003) that this joint ML solution on $r(x, y)$ and $\{\mu_j\}$ is identical to the ML solution on the sole $\{\mu_j\}$ assuming a gaussian prior probability on $r(x, y)$. The preliminary results show that, as in the previous method (estimation of the shift between two images) the criterion which considers the real noise variance contains a lot of local minima. This induce low performance of the method in this case. But, if we consider a constant noise variance, and we use the shift estimated with the previous method as guess for the minimization of this joint criterion, the performance are better in the low photon level domain (10 to 100).

References

- Blanc, A. & Mugnier, L. M. & Idier, J. 2003, JOSA A, 20, 6
- Carfantan, H. & Rougé, B. 2001, GRETSI XVIII, 849
- Downie & Walkup 1994, JOSA A, 11, 1599
- Gratadour, D. & Rouan, D. & Clenet, Y. & Gendron, E. & Lacombe, F. 2003 IAU Symp. 221, 265
- Groupe des Problèmes Inverses, "GPAV une grande oeuvre collective" internal report, LSS, 1997
- Guillaume, M. & Melon, P. & Refregier, P. & Llebaria, A. 1998, JOSA A, 15, 2841
- Slocumb, B. & Snyder, D. 1990, Acqu. Track. and Point. IV, 1304, 165
- Vander Lugt, A. 1964, IEEE Transactions on information theory, 10, 139
- Vijaya Kumar, B. & Dickey F. & DeLaurentis J. 1992, JOSA A, 9, 678